



Executive Headteacher: Mr C Reed

### Should your child be at school?

#### Ailments:

Coughs and colds	Your child should still attend school unless they have a raised temperature, shivers or drowsiness, in which case they should visit your GP and return to school 24 hours after they start to feel better. If your child has a long lasting cough, consult your GP and they will give guidance on school attendance.
Raised temperature	They should not attend school. They should return 24 hours after they feel better.
Headache	If your child has a minor headache they should attend school. If the headache is severe or accompanied by other symptoms such as drowsiness or vomiting, they should not attend and your GP should be consulted.
Vomiting and diarrhoea	They should not attend school until 48 hours after the last bout.
Bacillary dysentery	They should not attend school until physically fit and they have had no diarrhoea for 48 hours.
Sore throat	Your child should still attend school unless it is accompanied by a raised temperature.
Rashes	Rashes can be the sign of an infectious illness. If your child has a rash please check with your GP before they are sent into school. Please see the table below for further guidance.
Verrucae	They should come to school but keep foot covered at all times.
Conjunctivitis	Your child should come into school unless they cannot stop rubbing or touching their eyes.
Mumps	They should not attend school until the swelling has subsided, this would be 7 days minimum.
Hepatitis 'A'	Your child should not attend school until 7 days from the onset of jaundice.
Glandular fever	Your child should not attend school until advised by your GP that they are no longer infectious.
Head lice	Your child should be treated thoroughly before they can attend school.

Please note this guidance is separate to guidance surrounding COVID-19 symptoms.

#### Rashes:

Chickenpox	Stay at home until the last spot has scabbed over. This is usually 5 days from the onset of the rash.
German measles	Stay at home for 7 days from the onset of the rash.
Impetigo	Stay at home until the lesions are crusted or healed.
Measles	Stay at home for 4 days from the onset of the rash.
Scabies	Stay at home until the treatment is completed.
Scarlett fever	Stay at home until treatments have been given and your child is clinically recovered.

If you are unsure whether your child should come in to school please do contact the school or your doctor for advice.

